

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. Chan, M. Jakoobi, Y. Tian, C. Wang, N. Halcovitch, R. Boulatov and A. Sergeev, *Chem. Commun.*, 2021, DOI: 10.1039/D1CC03445F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

Selective, Radical-Free Activation of Benzylic C-H Bonds in Methylarenes

Received 00th January 20xx,
Accepted 00th January 20xxAntony P. Y. Chan,^a Martin Jakoobi,^a Chenxu Wang,^a Yancong Tian,^a Nathan Halcovitch,^b Roman Boulatov^a and Alexey G. Sergeev^{*a}

DOI: 10.1039/x0xx00000x


We report rare examples of exclusive benzylic C-H oxidative addition in industrially important methylarenes using simple η^4 -arene iridium complexes. Mechanistic studies showed that coordinatively unsaturated η^2 -arene intermediates are responsible for the selective activation of benzylic, not aromatic C-H bonds and formation of stable benzyl complexes after trapping with a phosphine ligand.

Syntheses of many fine and speciality chemicals rely on the C-H functionalisation of alkylarenes.¹ However, such C-H functionalisation without directing groups occurs unselectively and yields mixtures of products. Substantial progress has recently been made in selective functionalisation of unactivated arene Csp²-H bonds using transition metal catalysts,^{2,3} but the selective activation of benzylic Csp³-H bonds remains challenging^{1,4} because the competing activation of the aromatic C-H bonds is generally thermodynamically and kinetically preferred (Scheme 1A).⁵⁻⁸ Known examples of catalytic functionalisation of benzylic C-H bonds typically involve radical intermediates and stoichiometric amounts of peroxides, or strong bases, which limit the synthetic utility of the methods.^{1,4} Catalytic activation of benzylic C-H bonds by metal complexes in the absence of radical or basic reagents occurs much less common and often suffers from selectivity issues.⁹⁻¹⁵

The key step for designing general and selective catalytic benzylic C-H functionalisation is selective activation, e. g. oxidative addition of a benzylic C-H bond in a range of unactivated alkylarenes. However, only few complexes mediate oxidative addition of benzylic C-H bonds and, in most cases, is accompanied by oxidative addition of aromatic C-H bonds.^{16,17} The scope for each system is limited to one methylarene with sterically hindered arene C-H bonds such as mesitylene (1,3,5-trimethylbenzene); other arenes give mainly aromatic C-H activation products. Hence, the problem of highly selective benzylic C-H activation remains to be addressed.

Here we report the first general metal system that mediates exclusive benzylic C-H activation in nine unactivated arenes bearing from one to five methyl groups in the aromatic ring (Scheme 1B). Our kinetic and computational studies suggest that the key to the unique selectivity is generation of coordinatively unsaturated η^2 -arene intermediate, which preferentially activates benzylic, not aromatic C-H bonds in methyl arenes.

A Typical C-H activation in unactivated arenes



B This work: exclusive benzylic C-H activation




Figure 1. Metal-mediated C-H activation (oxidative addition) by metal complexes.

During our investigation of arene C-C cleavage in rare Cp*Ir(η^4 -arene) complexes,^{18,19} we found that thermolysis of the mesitylene complex in the presence of PMe₃ does not lead to C-C cleavage, but cleanly affords a product of benzylic C-H activation, Cp*Ir(CH₂C₆H₃Me_{2-3,5})(PMe₃).²⁰ This selective benzylic C-H activation, however, could be explained by the steric hindrance of aromatic C-H bonds. To test if benzylic C-H activation would occur in the least sterically hindered methylarene, toluene, we heated the iridium complex Cp*Ir(η^4 -toluene) (**1**) in the presence of 4 equiv. of PMe₃ in *n*-hexane at 100 °C for 1 h. Remarkably, all three isomers of the starting complex smoothly converted into a single benzylic C-H activation product (**2**) in a 97% isolated yield (Fig. 2A). This is the first example of exclusive benzylic C-H oxidative addition in an unhindered methylarene. Nearly quantitative yields of **2** were also obtained upon thermolysis of **1** in the presence of 4 equiv. of more sterically hindered phosphines such as PPh₃ and PCy₃ (Fig. 2A).²¹ PMe₃ was further used as the phosphine choice because the corresponding product **2** can be more conveniently isolated by the simple removal of the excess of the volatile phosphine *in vacuo*.

^a Department of Chemistry, University of Liverpool, Crown Street, Liverpool, L69 7ZD, United Kingdom. E-mail: sergeev@liverpool.ac.uk

^b Department of Chemistry Lancaster University Bailrigg, Lancaster, LA1 4YW, United Kingdom.

[†]Electronic Supplementary Information (ESI) available: CCDC 2090576 (**2b**). For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/x0xx00000x.




Figure 2. A) Iridium-mediated selective benzylic C-H activation of toluene upon thermolysis of **1** in the presence of phosphines. B) Crystal structure of **2b** with displacement ellipsoids drawn at 50% probability level except for H atoms. Hydride ligand was modelled for clarity. [a] ¹H NMR yield.

The structure of **2b** was confirmed by X-ray diffraction (Fig. 2A) and the hydride ligand, which was not explicitly detected by XRD, was observed in the ^1H NMR spectrum as a doublet at -16.4 ppm.

Building on the successful result with toluene (Fig. 2), we explored the scope of benzylic C-H activation in a range of methylarenes bearing from two to five methyl groups (Fig. 3). The thermolysis of the corresponding η^4 -arene complexes in *n*-hexane at 100 °C for 1-48 h gave C-H oxidative addition products **3-10** in high yields (84-93%). As shown in Fig. 3, the reactivity of the coordinated arenes strongly depends on the accessibility of methyl groups: arenes with unhindered methyl groups fully reacted within 1 h (**2a**, **4** and **5**), with methyl groups having one or two neighbouring substituents - 2 h (**3**, **8** and **9**) and 3 h (**7**), respectively. Ultimately, the pentamethylbenzene complex [Cp*Ir (η^4 -C₆Me₅H)] was consumed within 48 h (**10**), while its fully substituted congener was the least reactive and reacted to only 45% conversion after 168 h.²²




Figure 3. Scope of iridium-mediated benzylic C-H activation in coordinated methyl arenes. All yields reported are isolated yields. [a] Ratio of regioisomers.

Remarkably, the C-H activation occurred with high chemoselectivity, affording exclusively benzylic complexes in eight examples out of nine (Fig. 3). The only exception was the activation of *p*-xylene (**5**), which gave a 69:31 ratio of benzylic to aromatic C-H oxidative addition products.²³ To the best of our knowledge, this is the most general and selective system for oxidative addition of benzylic C-H bonds in alkyl arenes.

Another remarkable aspect of this benzylic C-H activation is that the regioselectivity does not appear to depend on steric effects. In case of 1,2,3-trimethylbenzene, **7**, (Fig. 3), the regioselectivity for activation of either of the two less hindered CH₃ groups vs the more hindered internal CH₃ equals 2.1:1, which after statistical correction for the number of methyl groups gives nearly equal regioselectivity of 1.1:1. Similarly, in pentamethylbenzene all three types of methyl groups have similar reactivity as evidenced by the statistically adjusted regioselectivity of 1.3:0.9:1 (**10**). More surprisingly, in 1,2,4-trimethylbenzene either of the more hindered CH₃ groups is greater than two times more reactive than the sterically unhindered CH₃ group.²⁴ This regioselectivity contrasts to known C-H activations in alkylarenes, where the least sterically hindered aromatic C-H bonds are cleaved preferentially.

To get insight into the mechanism of the reaction, we first conducted kinetic studies using thermolysis of mesitylene complex $\text{Cp}^*\text{Ir}(\eta^4\text{-mesitylene})$ (**11**) and PMe_3 (Fig. 4A). The reason for this choice is that **11** and product **6** exist as single isomers, which simplify monitoring and interpretation of the data. The thermolysis was conducted in cyclohexane- d_{12} at 75 °C and monitored by ^1H NMR. We measured the initial rates of the reaction by varying the concentration of PMe_3 and **11** independently. The rate of the reaction was first order in [**11**] and zero-order in $[\text{PMe}_3]$ (Figs. S3 and S5). Analogous tentative data were obtained for PPh_3 and PCy_3 . Note that although the variation of the concentration of these bulky phosphines did not change the rate of the reaction, the use of 1 equiv. instead of 4 equiv. of the phosphine relative to **11** led to the formation of the C-C cleavage side products.²⁵




Figure 4. (A) The model reaction to determine reaction order in Cp*Ir(η^4 -mesitylene) (**11**) and PMe₃. (B) Thermolysis of **11**-d₉ and **11**-d₃: kinetic isotope effects and the lack of H/D scrambling. [a] KIEs were calculated from the initial rates for separate reactions of the deuterated and non-deuterated species.

The observed preferential activation of weaker benzylic C-H bonds may suggest the involvement of radical pathways.⁴ One of the common tests for the participation of the radical mechanisms is the addition of radical traps such as TEMPO that inhibits radical reactions.²⁶ We probed this scenario by conducting thermolysis of **11** in the presence of 1 equiv. of TEMPO at 100 °C for 1 h. Full consumption of **1** was observed and the expected hydride product **6** was obtained in a 14% yield, which suggests that the involvement of radical intermediates is unlikely. The lower yield of **6** in the presence of the radical trap as compared to that in its absence (91%, Fig. 3) might be explained by the TEMPO-induced decomposition of **6** under reaction conditions as confirmed by control experiments.^{27,28} The lack of involvement of radicals in the benzylic C-H oxidative addition also agrees with the absence of observable deuterium incorporation of deuterium from the solvent upon thermolysis of **6** in d₁₂-cyclohexane (Fig. 4A).

To confirm that the hydride ligand in **6** results from the oxidative addition of the methyl groups of the arene ligand, we heated the model complex **11-d₉** having fully deuterated methyl groups on the aryl ligand (Fig. 4B). This led to the exclusive formation of the deuteride product **6-d₉**. Heating of the model complex **11-d₃** bearing deuterium labels on the three aromatic positions predictably gave hydride complex **6-d₃**. No observable H/D scrambling was detected in either case. Kinetic isotope effect measurements conducted in separate experiments with **11-d₃** and **11-d₉** complexes and the nondeuterated **11** resulted in negligible KIEs ($k_H/k_D = 1.03 \pm 0.18$ and 0.99 ± 0.09), which suggests that the rate-determining step does not involve benzylic or aromatic C-H bond cleavage (Figs. S7 and S9). To identify the mechanism that agrees with the observed experimental data and explains the unusual chemoselectivity, we computed a range of energy pathways for benzylic and

aromatic C-H activation in **6**. We conducted all geometry optimizations, reaction path calculations and calculations of thermodynamic corrections with the B3LYP functional and a mixed basis set of LANL2DZ for Ir and 6-31G(d) for all other atoms, which gave excellent results for the calculations of C-C oxidative addition in η^4 -arene iridium complexes.²⁰

The DFT computations indicated that the benzylic C-H activation **11**→**6** (Fig. 5, green path) has indeed the lowest energy as compared to the aromatic C-H activation **11**→**17** (Fig. 5, red and blue paths). The benzylic C-H oxidative addition **11**→**6** involves the initial rate-determining η^4 to η^2 arene ring slippage in the starting complex over the barrier of 24.4 kcal/mol. The resulting η^2 -arene intermediate **12** undergoes facile C-H activation of a methyl group to form an η^3 -benzyl complex **13** that is subsequently trapped by PMe₃ to yield the isolated benzylic complex **6**. The benzylic C-H activation preferentially occurs via the phosphine-free η^2 -complex **12**, rather than via phosphine ligated complex **14** because the latter is much less kinetically accessible from the starting complex **11** either directly or via **12** (Fig. 5).

All paths for aromatic C-H activation **11**→**17** also involve the η^2 -arene complex **12** as an essential intermediate. The direct aromatic C-H oxidative addition **12**→**16** requires a 20.1 kcal/mol higher barrier than the corresponding benzylic C-H activation **12**→**13** (Fig. 5; compare red and green paths). A much more favourable path to **17** involves the initial benzylic C-H activation **12**→**13** followed by the aromatic C-H activation yielding the dihydride intermediate **15**. The dihydride **15** reforms the benzylic C-H bond and the resulting coordinatively unsaturated aryl hydride **16** picks up PMe₃ to give the final product **17**. This lowest energy aromatic C-H oxidative addition still has a 6.6 kcal/mol higher barrier than the benzylic C-H activation path (Fig. 5, compare **13**→**16** vs **13**→**6**).




Figure 5. The calculated mechanisms for benzylic and aromatic C-H activation in **11** in the presence of added PMe₃ ligand. Energies at M06-L/(6-311+G*+LANL2TZ/cpcm=hexane)//B3LYP/(6-31G*+LANL2DZ) and thermodynamic corrections at 50 °C.

The results of DFT calculations indicate that the generation of a phosphine-free $\text{Cp}^*\text{Ir}(\eta^2\text{-arene})$ intermediate is the key to the exclusive regioselectivity of benzylic C-H activation in $\eta^4\text{-arene}$ iridium complexes (Fig. 6A). That is, the Cp^*Ir fragment selectively activates benzylic C-H bonds in the $\eta^2\text{-methyl arene}$ ligand and then picks up a phosphine ligand to afford the final product (Fig. 6A). The reactivity of this Cp^*Ir fragment is orthogonal to the reactivity of the corresponding well-known phosphine-ligated species, $\text{Cp}^*\text{Ir}(\text{PMe}_3)_3$, which preferentially activates aromatic C-H bonds as reported by Jones²⁹ and Bergman (Fig. 6B).³⁰

These $\text{Cp}^*\text{Ir}(\eta^2\text{-arene})$ intermediates have unique reactivity not only in the benzylic C-H activation. As we showed previously, when these species are generated in the absence of an added ligand, they promote oxidative addition of the aromatic C-C bonds.^{7,8} The starting $\text{Cp}^*\text{Ir}(\eta^4\text{-arene})$ complexes serve as excellent precursors for the generation of these highly reactive intermediates.

In summary, we reported an unprecedented highly selective oxidative addition of benzylic C-H bonds in a range of methylarenes, which occurs in simple $\text{Cp}^*\text{Ir}(\eta^4\text{-arene})$ complexes. Experimental and computational data suggest that the process occurs via the rate-determining formation of highly reactive $\eta^2\text{-arene}$ iridium species that undergo selective oxidative addition of benzylic C-H bonds. The unique reactivity of the $\eta^2\text{-arene}$ iridium species in benzylic C-H activation provides an exciting starting point for the design of novel highly selective functionalisations, which are currently investigated in our laboratory.

We thank the Leverhulme Trust, UK (Grant RPG-2018-406), EPSRC Early Career Fellowship (EP/L000075/1 to R.B.) and Petroleum Research Fund (58885-ND7 to R.B.) for financial support. Computations reported here were performed in the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation grant number ACI-1548562, with computational resources provided by the SDSC under allocation TG-CHE140039. Analytical services were provided by University of Liverpool.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- R. Vanjari and K. N. Singh, *Chem. Soc. Rev.*, 2015, **44**, 8062-8096.
- P. Wedi and M. van Gemmeren, *Angew. Chem. Int. Ed.*, 2018, **57**, 13016-13027.
- N. Kuhl, M. N. Hopkinson, J. Wencel-Delord and F. Glorius, *Angew. Chem. Int. Ed. Engl.*, 2012, **51**, 10236-10254.
- R. Yazaki and T. Ohshima, *Tetrahedron Lett.*, 2019, **60**, 151225.
- W. D. Jones and F. J. Feher, *J. Am. Chem. Soc.*, 1982, **104**, 4240-4242.
- P. E. M. Siegbahn, *J. Phys. Chem.*, 1995, **99**, 12723-12729.
- W. D. Jones and E. T. Hessell, *J. Am. Chem. Soc.*, 1993, **115**, 554-562.
- E. Clot, C. Mégret, O. Eisenstein and R. N. Perutz, *J. Am. Chem. Soc.*, 2006, **128**, 8350-8357. DOI: 10.1039/D1CC03445F
- S. Shimada, A. S. Batsanov, J. A. K. Howard and T. B. Marder, *Angew. Chem. Int. Ed.*, 2001, **40**, 2168-2171.
- T. Ishiyama, K. Ishida, J. Takagi and N. Miyaura, *Chem. Lett.*, 2001, **30**, 1082-1083.
- M. A. Larsen, C. V. Wilson and J. F. Hartwig, *J. Am. Chem. Soc.*, 2015, **137**, 8633-8643.
- W. N. Palmer, J. V. Obligacion, I. Pappas and P. J. Chirik, *J. Am. Chem. Soc.*, 2016, **138**, 766-769.
- C. M. Kelly, J. T. Fuller, C. M. Macaulay, R. McDonald, M. J. Ferguson, S. M. Bischof, O. L. Sydora, D. H. Ess, M. Stradiotto and L. Turculet, *Angew. Chem. Int. Ed.*, 2017, **56**, 6312-6316.
- T. Furukawa, M. Tobisu and N. Chatani, *Chem. Commun.*, 2015, **51**, 6508-6511.
- K. Mertins, A. Zapf and M. Beller, *J. Mol. Catal. A: Chem.*, 2004, **207**, 21-25.
- Y. Zhu, L. Fan, C.-H. Chen, S. R. Finnell, B. M. Foxman and O. V. Ozerov, *Organometallics*, 2007, **26**, 6701-6703.
- P. Burger and R. G. Bergman, *J. Am. Chem. Soc.*, 1993, **115**, 10462-10463.
- M. Jakoobi, N. Halcovitch, G. F. S. Whitehead and A. G. Sergeev, *Angew. Chem. Int. Ed.*, 2017, **56**, 3266-3269.
- M. Jakoobi, Y. Tian, R. Boulatov and A. G. Sergeev, *J. Am. Chem. Soc.*, 2019, **141**, 6048-6053.
- Y. Tian, M. Jakoobi, R. Boulatov and A. G. Sergeev, *Chem. Sci.*, 2021, **12**, 3568-3579.
- Note that the reaction also proceeds smoothly with just one equivalent of PMe_3 but C-C cleavage is not fully suppressed in the case of the larger phosphines.
- Heating $\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{Me}_6)$ and PMe_3 at 150 °C for 48 h led to the full conversion of the starting complex but gave a complex mixture of products. See the Supporting Information.
- This selectivity is reverse to that of the $\text{Cp}^*\text{Ir}(\text{PMe}_3)_3$ species generated *in situ* from $\text{Cp}^*\text{Ir}(\text{PMe}_3)_3\text{H}_2$, which reacts with *p*-xylene to afford the corresponding benzylic and aromatic C-H activated products in a 29:71 ratio, respectively. See Ref. 30.
- Although the origin of this selectivity has yet to be investigated, the selectivity results from kinetic, not thermodynamic control as evidenced by the DFT calculations of the relative stability of the corresponding products, which scale as **8c** > **8a** > **8b** (Fig. S11).
- See the Supporting Information.
- K. Matyjaszewski, *Macromolecules*, 1998, **31**, 4710-4717.
- A. C. Albeniz, P. Espinet, R. Lopez-Fernandez and A. Sen, *J. Am. Chem. Soc.*, 2002, **124**, 11278-11279.
- Heating **6** in the presence of 1 equiv. of TEMPO at 100 °C for 1 h led to 83% degradation of **6**. Please see the supporting information. TEMPO-induced decomposition of related palladium benzyl hydride complexes is described in Ref 27.
- W. D. Jones and F. J. Feher, *J. Am. Chem. Soc.*, 1984, **106**, 1650-1663.
- A. H. Janowicz and R. G. Bergman, *J. Am. Chem. Soc.*, 1983, **105**, 3929-3939.